

INCOMPRESSIBLE EULER EQUATIONS AND THE EFFECT OF CHANGES AT A DISTANCE

ELAINE COZZI AND JAMES P. KELLIHER

ABSTRACT. Because pressure is determined globally for the incompressible Euler equations, a localized change to the initial velocity will have an immediate effect throughout space. For solutions to be physically meaningful, one would expect such effects to decrease with distance from the localized change, giving the solutions a type of stability. Indeed, this is the case for solutions having spatial decay, as can be easily shown. We consider the more difficult case of solutions lacking spatial decay, and show that such stability still holds, albeit in a somewhat weaker form.

1. INTRODUCTION AND STATEMENT OF THE MAIN THEOREM

The Euler equations governing incompressible inviscid fluid flow in \mathbb{R}^d are given by

$$(E_d) \quad \begin{cases} \partial_t u + u \cdot \nabla u = -\nabla p \\ \operatorname{div} u = 0 \\ u|_{t=0} = u^0. \end{cases}$$

There is extensive literature on existence and uniqueness of solutions to the Euler equations on \mathbb{R}^d in various function spaces. Short time existence of smooth solutions to (E_d) goes back to papers of Lichtenstein and Gunther [12, 9]; [10] contains a modern form of these results. For (E_2) , surveys on global in time existence of solutions can be found in [13] and [4]. A classical result for weak solutions of the Euler equations in the plane is due to Yudovich [20], which gives existence and uniqueness of solutions with velocity in $L^2(\mathbb{R}^2)$ and vorticity (curl of the velocity) in $L^p \cap L^\infty(\mathbb{R}^2)$ for some $p < \infty$. (Yudovich worked in a bounded domain, but his argument is easily extended to the full plane.) In recent years, some attention has also been paid to the Euler equations with initial vorticity which does not decay at infinity or may be unbounded (see, for example, [1], [7], [15], [16], [17], [18], [21]).

We consider the question of what happens to the solution to the Euler Equations in \mathbb{R}^d at time $t > 0$ when a change is made to the initial velocity far away from a given point. We focus, in particular, on solutions to the Euler equations in two or three dimensions which have bounded vorticity and bounded velocity, but for which vorticity has no spatial decay.

For solutions having spatial decay of vorticity, bounds on the effects of far-away changes can be obtained in an elementary manner in the L^2 -norm of the velocity. We quickly review the argument, as it helps motivate what follows.

Date: July 31, 2015.

2010 Mathematics Subject Classification. Primary 76B03.

Key words and phrases. Fluid mechanics, Euler equations.

Write $B_R(x)$ for the ball of radius $R > 0$ centered at $x \in \mathbb{R}^d$. Let $\varphi \in C_C^\infty(\mathbb{R}^d)$ be supported in $B_2(0)$ with $\varphi \equiv 1$ on $B_1(0)$, and define $\varphi_R(x) = \varphi(x/R)$. Let (u_1, p_1) , (u_2, p_2) be two solutions to the Euler equations, and let $w = u_1 - u_2$, $p = p_1 - p_2$. Then subtracting (E_d) for u_2 from (E_d) for u_1 , multiplying by $\varphi_R^2 w$, and integrating over time and space, one obtains

$$(1.1) \quad \|\varphi_R w(t)\|_{L^2}^2 = \|\varphi_R w^0\|_{L^2}^2 + \int_0^t \int_{\mathbb{R}^d} \left[2(\varphi_R \nabla \varphi_R \cdot u_1) |w|^2 - 2(\varphi_R w \cdot \nabla u_2) \cdot (\varphi_R w) + 4p \varphi_R \nabla \varphi_R \cdot w \right].$$

For simplicity, suppose $\nabla u_2 \in L^\infty([0, t] \times \mathbb{R}^d)$. Then, assuming that our solutions have sufficient decay to ensure that

$$A := \|\nabla u_2\|_{L^\infty([0, t] \times \mathbb{R}^d)} + \sum_{j=1}^2 \left[\|u_j\|_{L^\infty([0, t] \times \mathbb{R}^d)} + \|u_j\|_{L^\infty(0, t; L^2)} + \|p_j\|_{L^\infty(0, t; L^2)} \right] < \infty,$$

we have

$$\|\varphi_R w(t)\|_{L^2}^2 \leq \|\varphi_R w^0\|_{L^2}^2 + \int_0^t \left[\frac{C}{R} A^3 + 2A \|\varphi_R w(s)\|_{L^2}^2 + \frac{4A^2}{R} \right] ds.$$

Gronwall's lemma then gives the bound,

$$(1.2) \quad \|\varphi_R w(t)\|_{L^2}^2 \leq \left[\|\varphi_R w^0\|_{L^2}^2 + \frac{C \max\{1, A\}^3}{R} t \right] e^{2At}.$$

The bound in (1.2) controls how rapidly the values of the initial velocity outside of $B_{2R}(0)$ can affect the value of the velocity inside $B_R(0)$, showing that, for decaying solutions, the Euler equations have a degree of locality. (Of course, many variations on this theme are possible.)

If we only have bounded, but decaying vorticity, say vorticity lying in $L^\infty([0, t]; L^1 \cap L^\infty)$, then we cannot conclude that ∇u_2 is bounded. Regardless, an argument like that in [21], employing the Calderon-Zygmund inequality, can be used to handle the integral involving ∇u_2 in (1.1). Osgood's lemma is used in place of Gronwall's lemma, leading to a different bound, but we suppress these details in this brief overview.

Mere membership of u_1, u_2 in $L^\infty([0, t]; L^2)$ gives that $\|\varphi_R w(t)\|_{L^2} \rightarrow 0$ as $R \rightarrow \infty$; it does not, however, give a rate of convergence. The rate of convergence in (1.2) comes from bounding the rate of decay of the "tail" of the velocity, and through it the pressure.

If we now assume that the velocity and vorticity are bounded but do not decay at infinity, the type of argument that led to (1.2) fails completely. The lack of decay introduces several difficulties when estimating terms in (1.1). First, while it is known that in 2D, $\nabla p \in L^\infty([0, t] \times \mathbb{R}^2)$ and $|p(t, x)| \leq C(t) \log(|x| + e)$ ([15, 11]), this information is insufficient to control the pressure term in (1.1). Second, the integral involving u_1 in (1.1) can no longer be controlled since $w \notin L^2(\mathbb{R}^d)$. Third, the integral involving ∇u_2 cannot be controlled using the approach of Yudovich in [21], as the vorticity only lies in $L^\infty(\mathbb{R}^d)$, and the Calderon-Zygmund theory does not apply in this setting. Finally, we would expect $\|\varphi_R w(t)\|_{L^2}^2$ to grow with R anyway; hence, it would appear that such

an estimate is unattainable when the solutions do not decay. We should seek instead a weighted L^∞ -based estimate.

One can view the bound in (1.2) as a type of stability estimate that demonstrates a partial locality of the solutions. Without some such locality, the very physical meaning of the solutions must be called into question: If a small change to the initial velocity at an arbitrarily large distance from a given point can immediately and significantly change the value of the solution at that point, then one could never effectively compute a solution.

The original motivation for this paper was to address this issue of physical meaning for the case of bounded vorticity, bounded velocity (Serfati) solutions to the 2D Euler equations. Such solutions were first derived by Serfati in [15], where their existence and uniqueness was obtained. Serfati's uniqueness argument easily extends to stability estimates in the L^∞ -norm of the velocity, but that argument cannot be localized. Locality of Serfati solutions has been an unaddressed problem.

Stability estimates in L^∞ -based norms for Serfati solutions were established in [1, 17]. The estimate obtained in [1] is an outgrowth of Serfati's uniqueness proof in [15], while the estimate in [17] is an outgrowth of Vishik's uniqueness proof in [18], and both [15] and [18] depend critically on L^∞ estimates. (Neither employs an energy argument, nor is the maximum principle a key ingredient.) Our approach, which we describe in more detail below, will be to introduce a weighted L^∞ -norm into Vishik's argument, adapting as well some pressure-based estimates in [17].

The weight we use is defined as follows. For fixed $z \in \mathbb{R}^d$ and $\alpha > 0$, define $g_z : \mathbb{R}^d \rightarrow \mathbb{R}$ by

$$g_z(x) = (1 + |x - z|)^{-\alpha}.$$

A higher value of α makes the weighted norm more localized. As one might expect, the behavior of the Eulerian pressure limits how large α can be. In particular, the important properties of the Riesz operators described in Theorem 7 below play a crucial role in determining this upper limit.

For a velocity field, u , on \mathbb{R}^d , we define the vorticity, $\omega = \omega(u)$, as

$$\omega = \nabla u - (\nabla u)^T.$$

(In 2D, one can alternately use $\omega = \partial_1 u^2 - \partial_2 u^1$.)

We now state our main result.

Theorem 1. *For fixed $T > 0$, assume that (u_1, p_1) and (u_2, p_2) satisfy (E_d) in $\mathcal{D}'((0, T) \times \mathbb{R}^d)$, with*

$$\begin{aligned} u_1, u_2 &\in L^\infty([0, T]; L^p \cap L^\infty(\mathbb{R}^d)) \text{ for } p \in [1, \infty], \\ \omega_1, \omega_2 &\in L^\infty([0, T]; L^\infty(\mathbb{R}^d)), \\ p_1 &= \sum_{k,l=1}^d R_k R_l u_1^k u_1^l, \quad p_2 = \sum_{k,l=1}^d R_k R_l u_2^k u_2^l \text{ in BMO.} \end{aligned}$$

Let

$$M = M(t) := \sup_{0 \leq s \leq t} (1 + \|\omega_1(s)\|_{L^\infty} + \|\omega_2(s)\|_{L^\infty} + \|u_1(s)\|_{L^\infty \cap L^p} + \|u_2(s)\|_{L^\infty \cap L^p}),$$

and set $\epsilon_0 = \|g_z(u_1^0 - u_2^0)\|_{L^\infty}$. Define the strictly increasing function, $\mu: [0, 1] \mapsto [0, 1]$ by $\mu(x) = x(1 - \log x)$. Then, whenever $\alpha < 1 + d/p$,

$$\|g_z(u_1 - u_2)(t)\|_{L^\infty} \leq M\mu \left(\min \left\{ C \frac{\epsilon_0}{M} + CMt\mu \left(\left(\frac{\epsilon_0}{M} \right)^{e^{-CMt}} \right), 1 \right\} \right).$$

The constant C depends only upon d and α .

Remark 2. We can see from Theorem 1 that for any $t \leq T$, we obtain a useful bound on $\|g_z(u_1 - u_2)(t)\|_{L^\infty}$ as long $\|g_z(u_1^0 - u_2^0)\|_{L^\infty}$ is sufficiently small. Also, stability in $C^r(\mathbb{R}^d)$ for all $r < 1$ follows from Theorem 1 by interpolation.

We now briefly describe three examples to which Theorem 1 applies. All of these examples are 2D, for even though Theorem 1 is not dimension-dependent, only in 2D are weak solutions to the Euler equations known to exist.

If $p = \infty$ and $d = 2$, then Theorem 1 above addresses Serfati (bounded vorticity, bounded velocity) solutions to the Euler equations in the plane (see [15]), and supplies an alternate proof of uniqueness to those in [15, 1, 17]. Moreover, it is shown in [8] that for Serfati solutions,

$$\|u_j(t)\|_{L^\infty} \leq C\|u_j^0\|_{L^\infty} (1 + \|\omega_j^0\|_{L^\infty} t).$$

Hence, $M(t) \leq C(M(0) + M(0)^2 t)$ for Serfati solutions.

If u^0 belongs to $L^2(\mathbb{R}^2)$ and ω^0 belongs to $L^\infty(\mathbb{R}^2)$, then there exists a global-in-time solution to (E_2) which conserves energy and for which vorticity remains bounded (see [6] for details). For such solutions, one can apply a simple argument using Littlewood-Paley theory to show that for all $t > 0$,

$$\|u(t)\|_{L^\infty} \leq C(\|u(t)\|_{L^2} + \|\omega(t)\|_{L^\infty}) = C(\|u^0\|_{L^2} + \|\omega^0\|_{L^\infty}).$$

Thus, Theorem 1 applies for this class of solutions with $p = 2$ and $d = 2$, and we have $M(t) \leq CM(0)$.

If u^0 belongs to $L^2(\mathbb{R}^2)$ and ω^0 belongs to $L^p \cap L^\infty(\mathbb{R}^2)$ for some $p < \infty$, then Theorem 1 addresses the classical Yudovich solutions to (E_2) and we again have $M(t) \leq CM(0)$.

Let us consider one simple application of Theorem 1, say, to be explicit, in the setting of Serfati solutions. Suppose that $u_1^0 = u_2^0$ on $B_R(0)$ for some $R > 0$ and that u_1^0 is compactly supported on $B_R(0)$. Then

$$(1.3) \quad \epsilon_0 = \|g_0(u_1^0 - u_2^0)\|_{L^\infty} \leq \|u_2^0\|_{L^\infty(B_R(0))^c} (1 + R)^{-\alpha}.$$

The bound that results from Theorem 1 at time zero is

$$|(u_1^0 - u_2^0)(0)| \leq \|g_0(u_1^0 - u_2^0)\|_{L^\infty} \leq CM\mu(C\epsilon_0/M).$$

Because μ is superlinear ($\lim_{x \rightarrow 0^+} \mu'(x) = +\infty$), the bound at time zero is somewhat higher than the initial difference. Nonetheless, because the bound on ϵ_0 in (1.3) decreases inversely with R , the bound on $\|g_0(u_1 - u_2)(t)\|_{L^\infty}$ decreases as $R \rightarrow \infty$ for any fixed $t > 0$. This allows us to determine the distance, R , away from a given point (the origin, in this case) beyond which the initial velocity will not affect the value of the velocity at the given point up to a desired time.

This example illustrates that the change to the initial velocity in Theorem 1 need not be localized, it need only be far away from the point or points of interest. We see this for

spatially decaying solutions as well in (1.2), though there the control is stronger and can be more cleanly expressed.

Our proof of Theorem 1 is a localization of the uniqueness proof of Vishik's in [18], with the pressure, however, handled much as in [17]. We localize Vishik's argument using the function, g_z . Each of the terms that appear in [18] have an analog here, though most of these terms must be bounded in a different manner. At the heart of bounding these localized terms are the inequality,

$$(1.4) \quad g_z(x)h_z(x-y) \leq (1+|y|)^\alpha,$$

established in the proof of Lemma 6, and the identity,

$$(1.5) \quad \nabla g_z = \psi_z g_z,$$

where ψ_z is a vector-valued function with L^∞ -norm independent of $z \in \mathbb{R}^d$ (in fact, $\|\psi_z\|_{L^\infty} = \alpha$).

The identity in (1.4) is used several times, in each case to bound terms of the form $g_z(x)(\phi * f)(x)$, as follows:

$$\begin{aligned} g_z(x)(\phi * f)(x) &= \int \phi(y)g_z(x-y)g_z(x)h_z(x-y)f(x-y) dy \\ &\leq \int |\phi(y)|(1+|y|)^\alpha |(g_z f)(x-y)| dy = (|\phi(\cdot)|(1+|\cdot|)^\alpha * |g_z f|)(x). \end{aligned}$$

When ϕ has sufficient decay, this effectively brings g_z into the integral to obtain an estimate based on the function $g_z f$.

The identity in (1.5) is used to control the one term that is the analog of the integral involving u_1 in (1.1). As a consequence of (1.5), $|\nabla g_z| \leq C|g_z|$. Such a bound is never possible for a compactly supported function: this inability is the fundamental reason why we cannot use the "cleaner" localization that a compactly supported function would provide. This same issue shows up, for instance, in [8] (see Section 3.3), though it is addressed in a different manner.

In [18], decay of vorticity is assumed, which allows the pressure to be easily dealt with. Without such decay, the pressure requires much more careful treatment. We follow the approach in [17], localizing it, much as we localize the approach in [18].

Remark 3. Being a localization of the approaches in [18, 17], we expect that a result similar to Theorem 1 will hold for vorticity in the spaces containing unbounded functions ($bmo(\mathbb{R}^d)$, for example) considered in [18, 17]. However, for simplicity, we do not consider unbounded vorticity here.

Finally, we mention that the proof of uniqueness of Serfati solutions in [15] does not rely upon paradifferential calculus, and hence is not so tightly tied to the full plane. Indeed, Serfati's approach was extended to a 2D exterior domain in [1]. The proof involves bounding the difference between the flow maps for the two solutions (in this, it has similarities to the proof in [14] of uniqueness for Euler solutions in a 2D bounded domain). This approach seems resistant, however, to localization. It thus remains an interesting open problem whether an analog of Theorem 1 can be obtained for bounded vorticity, bounded velocity solutions in a 2D exterior domain.

2. DEFINITIONS AND PRELIMINARY LEMMAS

We first define the Littlewood-Paley operators. It is classical that there exists two functions $\chi, \phi \in S(\mathbb{R}^d)$ with $\text{supp } \hat{\chi} \subset \{\xi \in \mathbb{R}^d : |\xi| \leq \frac{5}{6}\}$ and $\text{supp } \hat{\phi} \subset \{\xi \in \mathbb{R}^d : \frac{3}{5} \leq |\xi| \leq \frac{5}{3}\}$, such that, if for every $j \geq 0$ we set $\phi_j(x) = 2^{jd}\phi(2^jx)$ then

$$\hat{\chi} + \sum_{j \geq 0} \hat{\phi}_j = \hat{\chi} + \sum_{j \geq 0} \hat{\phi}(2^{-j}\cdot) \equiv 1.$$

For $n \in \mathbb{Z}$ define $\chi_n \in S(\mathbb{R}^d)$ in terms of its Fourier transform $\hat{\chi}_n$, where $\hat{\chi}_n$ satisfies

$$\hat{\chi}_n(\xi) = \hat{\chi}(\xi) + \sum_{j \leq n} \hat{\phi}_j(\xi)$$

for all $\xi \in \mathbb{R}^d$. For $f \in S'(\mathbb{R}^d)$ define the operator S_n by

$$S_n f = \chi_n * f.$$

Finally, for $f \in S'(\mathbb{R}^d)$ and $j \geq -1$, define the Littlewood-Paley operators Δ_j by

$$\Delta_j f = \begin{cases} \chi * f, & j = -1 \\ \phi_j * f, & j \geq 0. \end{cases}$$

We will also need the paraproduct decomposition introduced by J.-M. Bony in [2]. We recall the definition of the paraproduct and remainder used in this decomposition.

Definition. Define the paraproduct of two functions f and g by

$$T_f g = \sum_{\substack{i, j \\ i \leq j-2}} \Delta_i f \Delta_j g = \sum_{j=1}^{\infty} S_{j-2} f \Delta_j g.$$

We use $R(f, g)$ to denote the remainder. $R(f, g)$ is given by the following bilinear operator:

$$R(f, g) = \sum_{\substack{i, j \\ |i-j| \leq 1}} \Delta_i f \Delta_j g.$$

Bony's decomposition then gives

$$fg = T_f g + T_g f + R(f, g).$$

We will make use of Bernstein's Lemma in what follows. A proof of the lemma can be found in [3], chapter 2.

Lemma 4. (*Bernstein's Lemma*) *Let r_1 and r_2 satisfy $0 < r_1 < r_2 < \infty$, and let p and q satisfy $1 \leq p \leq q \leq \infty$. There exists a positive constant C such that for every integer k , if u belongs to $L^p(\mathbb{R}^d)$, and $\text{supp } \hat{u} \subset B(0, r_1\lambda)$, then*

$$(2.1) \quad \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^q} \leq C^k \lambda^{k+d(\frac{1}{p}-\frac{1}{q})} \|u\|_{L^p}.$$

Furthermore, if $\text{supp } \hat{u} \subset C(0, r_1\lambda, r_2\lambda)$, then

$$(2.2) \quad C^{-k} \lambda^k \|u\|_{L^p} \leq \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^p} \leq C^k \lambda^k \|u\|_{L^p}.$$

We will also make use of Osgood's Lemma in the proof of Theorem 1. We refer the reader to [5] for a proof of the lemma.

Lemma 5. *Let ρ be a measurable positive function, let γ be a locally integrable positive function, and let μ be a continuous increasing function. Assume that for some $\beta > 0$, the function ρ satisfies*

$$\rho(t) \leq \beta + \int_{t_0}^t \gamma(s) \mu(\rho(s)) ds.$$

Then $-\phi(\rho(t)) + \phi(\beta) \leq \int_{t_0}^t \gamma(s) ds$, where $\phi(x) = \int_x^1 \frac{1}{\mu(r)} dr$.

In what follows, we set

$$h_z(x) = 1/g_z(x) = (1 + |x - z|)^\alpha.$$

The following lemma is a key technical ingredient in the proof of Theorem 1.

Lemma 6. *Assume f belongs to $L^\infty(\mathbb{R}^d)$. There exists a constant $C > 0$, depending only on α , such that for every $j \geq -1$,*

$$(2.3) \quad \|g_z \Delta_j f\|_{L^\infty} \leq C \|g_z f\|_{L^\infty},$$

and, for $1 \leq k \leq d$,

$$(2.4) \quad \|g_z \partial_k \Delta_j f\|_{L^\infty} \leq C 2^j \|g_z f\|_{L^\infty}.$$

Proof. We first prove (2.3). Observe that for any $\alpha > 0$,

$$(2.5) \quad g_z(x) h_z(x - y) = \left(\frac{1 + |x - y - z|}{1 + |x - z|} \right)^\alpha \leq \left(\frac{1 + |x - z| + |y|}{1 + |x - z|} \right)^\alpha \leq (1 + |y|)^\alpha.$$

Assume $j \geq 0$. By (2.5),

$$(2.6) \quad \begin{aligned} g_z(x) \int_{\mathbb{R}^d} \phi_j(y) f(x - y) dy &= g_z(x) \int_{\mathbb{R}^d} \phi_j(y) h_z(x - y) g_z(x - y) f(x - y) dy \\ &\leq C \|g_z f\|_{L^\infty} \int_{\mathbb{R}^d} |\phi_j(y)| g_z(x) h_z(x - y) dy \leq C \|g_z f\|_{L^\infty} \int_{\mathbb{R}^d} |\phi_j(y)| (1 + |y|)^\alpha dy \\ &\leq C \|g_z f\|_{L^\infty} \int_{\mathbb{R}^d} |\phi(y)| (1 + |2^{-j} y|)^\alpha dy \leq C \|g_z f\|_{L^\infty}. \end{aligned}$$

An identical argument can be applied for $j = -1$ with ϕ replaced by χ . This proves (2.3).

To establish (2.4), note that for $j \geq 0$,

$$(2.7) \quad \begin{aligned} g_z(x) \partial_k \Delta_j f(x) &= g_z(x) \int_{\mathbb{R}^d} \partial_k \phi_j(y) f(x - y) dy \\ &= 2^{j(d+1)} \int_{\mathbb{R}^d} (\partial_k \phi)(2^j y) g_z(x) h_z(x - y) g_z(x - y) f(x - y) dy \\ &\leq C 2^j \|g_z f\|_{L^\infty} \int_{\mathbb{R}^d} (\partial_k \phi)(y) (1 + |2^{-j} y|)^\alpha dy \leq C 2^j \|g_z f\|_{L^\infty}. \end{aligned}$$

Again, an identical argument can be applied for $j = -1$ with ϕ replaced by χ . This completes the proof of Lemma 6. \square

In the proof of Theorem 1, the main restriction on the decay of g_z , and hence on the value of α , comes from the pressure term. To estimate this term, we will need the following theorem from [19], which gives a pointwise bound on the Riesz operators applied to sufficiently decaying functions with moment conditions.

Theorem 7. *Assume f is a function on \mathbb{R}^d which satisfies:*

- 1) $|f(x)| \leq C(1 + |x|)^{-d-N+\epsilon}$,
- 2) $|D^\alpha f(x)| \leq C(1 + |x|)^{-d-N-1+\epsilon}$ for all $|\alpha| = 1$, and
- 3) $\int x^\beta f(x) dx = 0$ for all $|\beta| < N$

for some fixed $\epsilon \in [0, 1)$ and some positive integer N . Then for each i , $1 \leq i \leq d$,

$$|R_i f(x)| \leq C(1 + |x|)^{-d-N+\epsilon+\delta}$$

for every δ satisfying $0 < \delta < 1 - \epsilon$.

We will apply Theorem 7 to both $f(x) = \nabla \chi(x)$ and $f(x) = D^\beta \phi(x)$, where χ and ϕ are as above. In particular, we will make use of the following corollary to Theorem 7, which is proved in [7].

Lemma 8. *Assume η belongs to $\mathcal{S}(\mathbb{R}^d)$. Given $\epsilon > 0$, there exists $C > 0$ such that for each i, j with $1 \leq i, j \leq d$,*

$$|R_i R_j \nabla \eta(x)| \leq C(1 + |x|)^{-d-1+\epsilon}.$$

Moreover, if ψ belongs to $\mathcal{S}(\mathbb{R}^d)$ and has Fourier transform supported away from the origin, then given any $\gamma > 0$, there exists $C > 0$ such that for each i, j with $1 \leq i, j \leq d$,

$$|R_i R_j \psi(x)| \leq C(1 + |x|)^{-\gamma}.$$

We will also make frequent use of the following well-known lemma. A proof of the lemma can be found in [16].

Lemma 9. *Assume u is a divergence-free vector field in $L^2_{loc}(\mathbb{R}^d)$ with $\text{curl } u = \omega$. There exists $C > 0$ such that for every $j \geq 0$,*

$$(2.8) \quad \|\Delta_j \nabla u\|_{L^\infty} \leq C \|\Delta_j \omega\|_{L^\infty}.$$

3. PROOF OF THEOREM 1

The proof of Theorem 1 is a modification of the proof of uniqueness of solutions to the Euler equations, as given in [18, 17]. The proof below deviates from those in [18, 17] in that we must account for the weight g_z when proving the necessary estimates. To do this, we make use of Lemma 6 and Lemma 8.

Let u_1, u_2, g_z , and $T > 0$ be as in the statement of Theorem 1. Set $w = u_1 - u_2$. For any $N \geq 0$,

$$\begin{aligned}
 \|g_z w\|_{L^\infty} &\leq \sum_{j=-1}^N \|g_z \Delta_j w\|_{L^\infty} + \sum_{j>N} \|g_z \Delta_j w\|_{L^\infty} \\
 &\leq CN \sup_{-1 \leq j \leq N} \|g_z \Delta_j w\|_{L^\infty} + \sum_{j>N} \|\Delta_j w\|_{L^\infty} \\
 (3.1) \quad &\leq CN \sup_j \|g_z \Delta_j w\|_{L^\infty} + C \sum_{j>N} 2^{-j} (\|\Delta_j \omega_1\|_{L^\infty} + \|\Delta_j \omega_2\|_{L^\infty}) \\
 &\leq CN \sup_j \|g_z \Delta_j w\|_{L^\infty} + C \sum_{j>N} 2^{-j} (\|\Delta_j \omega_1\|_{L^\infty} + \|\Delta_j \omega_2\|_{L^\infty}) \\
 &\leq CN \sup_j \|g_z \Delta_j w\|_{L^\infty} + CM 2^{-N},
 \end{aligned}$$

where we used Bernstein's Lemma and Lemma 9 to get the third inequality. To estimate $\|g_z w\|_{L^\infty}$, we establish an estimate for $\sup_j \|g_z \Delta_j w\|_{L^\infty}$ and only then judiciously choose a value of N (see (3.28)).

We now estimate $\sup_j \|g_z \Delta_j w\|_{L^\infty}$, a task that will consume most of our effort.

Set $p = p_1 - p_2$. Note that w satisfies

$$\partial_t w + w \cdot \nabla u_2 + u_1 \cdot \nabla w = -\nabla p.$$

Applying the Δ_j operator to this equation gives

$$\partial_t \Delta_j w + S_{j-2} u_1 \cdot \nabla \Delta_j w = -Q_j(u_1, w) - \Delta_j(w \cdot \nabla u_2) - \nabla \Delta_j p,$$

where

$$(3.2) \quad Q_j(u_1, w) = \Delta_j(u_1 \cdot \nabla w) - S_{j-2} u_1 \cdot \nabla \Delta_j w.$$

We now multiply through by g_z to get

$$\partial_t(g_z \Delta_j w) + g_z S_{j-2} u_1 \cdot \nabla \Delta_j w = -g_z Q_j(u_1, w) - g_z \Delta_j(w \cdot \nabla u_2) - g_z \nabla \Delta_j p.$$

Observe that, by the product rule,

$$\begin{aligned} g_z S_{j-2} u_1 \cdot \nabla \Delta_j w &= S_{j-2} u_1 \cdot g_z \nabla \Delta_j w \\ &= S_{j-2} u_1 \cdot \nabla(g_z \Delta_j w) - (S_{j-2} u_1 \cdot \nabla g_z) \Delta_j w. \end{aligned}$$

This implies that

$$(3.3) \quad \partial_t(g_z \Delta_j w) + S_{j-2} u_1 \cdot \nabla(g_z \Delta_j w) = I + II + III + IV,$$

where

$$\begin{aligned} I &= (S_{j-2} u_1 \cdot \nabla g_z) \Delta_j w \\ II &= -g_z Q_j(u_1, w) \\ III &= -g_z \Delta_j(w \cdot \nabla u_2) \\ IV &= -g_z \nabla \Delta_j p. \end{aligned}$$

Fix $p_0 \geq 3$ to be specified later. We will estimate $I-IV$ for three cases: 1) middle frequencies, $3 \leq j \leq p_0$; 2) high frequencies, $j > p_0$; 3) low frequencies, $-1 \leq j \leq 2$, in that order.

So assume that $3 \leq j \leq p_0$. We begin with I . A simple calculation gives (1.5), which allows us to write

$$(3.4) \quad \begin{aligned} \|I\|_{L^\infty} &= \|(S_{j-2} u_1 \cdot \psi_z g_z) \Delta_j w\|_{L^\infty} \leq \alpha \|S_{j-2} u_1\|_{L^\infty} \|g_z \Delta_j w\|_{L^\infty} \\ &\leq CM \|g_z \Delta_j w\|_{L^\infty}. \end{aligned}$$

Here, we used $\|S_{j-2} u_1\|_{L^\infty} \leq C \|u_1\|_{L^\infty}$.

Next, we estimate III . Expanding III using Bony's paraproduct decomposition gives

$$(3.5) \quad \begin{aligned} III &= -g_z \Delta_j \sum_{|l-j| \leq 3} \Delta_l w \cdot \nabla S_{l-2} u_2 - g_z \Delta_j \sum_{|l-j| \leq 3} S_{l-2} w \cdot \nabla \Delta_l u_2 \\ &\quad - g_z \Delta_j \sum_{\substack{|l-l'| \leq 1, \\ \max\{l, l'\} \geq j-3}} \Delta_l w \cdot \nabla \Delta_{l'} u_2. \end{aligned}$$

The first term in (3.5) can be estimated using Lemma 6 as follows:

$$(3.6) \quad \begin{aligned} \left| g_z \Delta_j \sum_{|l-j| \leq 3} \Delta_l w \cdot \nabla S_{l-2} u_2 \right| &\leq C \sum_{|l-j| \leq 3} \|g_z \Delta_l w \cdot \nabla S_{l-2} u_2\|_{L^\infty} \\ &\leq C \sum_{|l-j| \leq 3} \|S_{l-2} \nabla u_2\|_{L^\infty} \|g_z \Delta_l w\|_{L^\infty} \leq CM(j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty}. \end{aligned}$$

To obtain the last inequality above, we applied Bernstein's lemma and Lemma 9 to establish the series of estimates,

$$(3.7) \quad \|S_{l-2} \nabla u_2\|_{L^\infty} \leq \|\Delta_{-1} \nabla u_2\|_{L^\infty} + \sum_{j=0}^{l-2} \|\Delta_j \nabla u_2\|_{L^\infty} \leq C \|u_2\|_{L^\infty} + C(l-2) \|\omega_2\|_{L^\infty}.$$

For the second term in (3.5), another application of Lemma 6 and Lemma 9 gives

$$(3.8) \quad \begin{aligned} \left| g_z \Delta_j \sum_{|l-j| \leq 3} S_{l-2} w \cdot \nabla \Delta_l u_2 \right| &\leq C \sum_{|l-j| \leq 3} \|g_z S_{l-2} w \cdot \nabla \Delta_l u_2\|_{L^\infty} \\ &\leq C \sum_{|l-j| \leq 3} \left(\|\Delta_l \nabla u_2\|_{L^\infty} \sum_{m \leq l-2} \|g_z \Delta_m w\|_{L^\infty} \right) \leq CM(j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty}. \end{aligned}$$

The third term in (3.5) can be estimated using the divergence-free property of w , Lemma 6, Bernstein's Lemma, and Lemma 9. We have the series of estimates

$$(3.9) \quad \begin{aligned} \left| g_z \Delta_j \sum_{\substack{|l-l'| \leq 1, \\ \max\{l, l'\} \geq j-3}} \Delta_l w \cdot \nabla \Delta_{l'} u_2 \right| &= \left| g_z \Delta_j \sum_{k=1}^d \sum_{\substack{|l-l'| \leq 1, \\ \max\{l, l'\} \geq j-3}} \partial_k (\Delta_l w^k \Delta_{l'} u_2) \right| \\ &\leq C \sum_{\substack{|l-l'| \leq 1, \\ \max\{l, l'\} \geq j-3}} 2^j \|g_z \Delta_l w \Delta_{l'} u_2\|_{L^\infty} \leq C \sum_{\substack{|l-l'| \leq 1, \\ \max\{l, l'\} \geq j-3}} 2^{j-l'} \|\Delta_{l'} \nabla u_2\|_{L^\infty} \|g_z \Delta_l w\|_{L^\infty} \\ &\leq C \|\omega_2\|_{L^\infty} \sup_q \|g_z \Delta_q w\|_{L^\infty} \leq CM \sup_q \|g_z \Delta_q w\|_{L^\infty}. \end{aligned}$$

Substituting (3.6), (3.8), and (3.9) into (3.5) yields the estimate

$$(3.10) \quad \|III\|_{L^\infty} \leq CM(j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty}.$$

We now estimate II . As in [18], we write $Q_j(u_1, w) = \sum_{m=1}^4 Q_j^m(u_1, w)$, where

$$\begin{aligned} Q_j^1(u_1, w) &= \sum_{k=1}^d \Delta_j T_{\partial_k w} u_1^k \\ Q_j^2(u_1, w) &= - \sum_{k=1}^d [T_{u_1^k} \partial_k, \Delta_j] w \\ Q_j^3(u_1, w) &= \sum_{k=1}^d T_{u_1^k - S_{j-2} u_1^k} \partial_k \Delta_j w \\ Q_j^4(u_1, w) &= \sum_{k=1}^d \{ \Delta_j R(u_1^k \partial_k w) - R(S_{j-2} u_1^k, \Delta_j \partial_k w) \}. \end{aligned}$$

To estimate $g_z Q_j^1(u_1, w)$, we use the divergence-free property of u_1 , Lemma 6, Bernstein's Lemma, and Lemma 9 to write

$$\begin{aligned} |g_z Q_j^1(u_1, w)| &= \left| g_z \sum_{k=1}^d \Delta_j \partial_k \sum_{|l-j| \leq 3} S_{l-2} w \Delta_l u_1^k \right| \leq C \sum_{|l-j| \leq 3} 2^j \|g_z S_{l-2} w \Delta_l u_1\|_{L^\infty} \\ &\leq C \sum_{|l-j| \leq 3} 2^{j-l} (l-2) \sup_q \|g_z \Delta_q w\|_{L^\infty} \|\Delta_l \nabla u_1\|_{L^\infty} \\ &\leq C \|\omega_1\|_{L^\infty} (j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty} \leq CM(j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty}. \end{aligned}$$

Next, we estimate $g_z Q_j^3(u_1, w)$. By properties of the Littlewood-Paley operators, there exists $M_0 > 0$ such that

$$\begin{aligned} Q_j^3(u_1, w) &= \sum_{k=1}^d \sum_{l=1}^{\infty} S_{l-2} (u_1^k - S_{j-2} u_1^k) \Delta_l \partial_k \Delta_j w \\ &= \sum_{k=1}^d \sum_{|l-j| \leq 3} S_{l-2} \left(\sum_{m=j-1}^{\infty} \Delta_m u_1^k \right) \Delta_l \partial_k \Delta_j w = \sum_{k=1}^d \sum_{|l-j| \leq 3} \left(\sum_{|m-j| \leq M_0} \Delta_m u_1^k \right) \Delta_l \partial_k \Delta_j w. \end{aligned}$$

Hence,

$$\begin{aligned} (3.11) \quad |g_z Q_j^3(u_1, w)| &\leq C \sum_{k=1}^d \sum_{|l-j| \leq 3} \left(\sum_{|m-j| \leq M_0} 2^{-m} \|\Delta_m \nabla u_1\|_{L^\infty} \right) \|g_z \partial_k \Delta_l \Delta_j w\|_{L^\infty} \\ &\leq C \|\omega_1\|_{L^\infty} \sum_{k=1}^d \sum_{|l-j| \leq 3} 2^{-j} \|g_z \partial_k \Delta_l \Delta_j w\|_{L^\infty}, \end{aligned}$$

where Bernstein's Lemma was used to get the first inequality and Lemma 9 was applied to get the second inequality. Now note that by Lemma 6,

$$\|g_z \partial_k \Delta_l \Delta_j w\|_{L^\infty} \leq C 2^l \|g_z \Delta_j w\|_{L^\infty}.$$

Inserting this estimate into (3.11) gives

$$|g_z Q_j^3(u_1, w)| \leq CM \sup_q \|g_z \Delta_q w\|_{L^\infty}.$$

To estimate $g_z Q_j^2(u_1, w)$, we observe that there exists $M_0 > 0$ such that

$$Q_j^2(u_1, w) = - \sum_{k=1}^d \sum_{|l-j| \leq M_0} [S_{l-2} u_1^k, \Delta_j] \partial_k \Delta_l w.$$

We express the commutator as a convolution integral and utilize the divergence-free property of u_1 to write,

$$\begin{aligned} Q_j^2(u_1, w)(x) &= - \sum_{|l-j| \leq M_0} \sum_{k=1}^d \int_{\mathbb{R}^d} \phi_j(y) (S_{l-2} u_1^k(x) - S_{l-2} u_1^k(x-y)) \partial_k \Delta_l w(x-y) dy \\ &= - \sum_{|l-j| \leq M_0} \sum_{k=1}^d \int_{\mathbb{R}^d} \partial_k \phi_j(y) (S_{l-2} u_1^k(x) - S_{l-2} u_1^k(x-y)) \Delta_l w(x-y) dy. \end{aligned}$$

Hence,

$$\begin{aligned} &|g_z Q_j^2(u_1, w)(x)| \\ &\leq g_z(x) \sum_{|l-j| \leq M_0} 2^{j(d+1)} \int_{\mathbb{R}^d} |\nabla \phi(2^j y)| \|S_{l-2} \nabla u_1\|_{L^\infty} |y| g_z(x-y) h_z(x-y) |\Delta_l w(x-y)| dy \\ &\leq \sum_{|l-j| \leq M_0} 2^{j(d+1)} \|g_z \Delta_l w\|_{L^\infty} \|S_{l-2} \nabla u_1\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla \phi(2^j y)| |y| g_z(x) h_z(x-y) dy \\ &\leq \sum_{|l-j| \leq M_0} 2^j \|g_z \Delta_l w\|_{L^\infty} \|S_{l-2} \nabla u_1\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla \phi(y)| |2^{-j} y| (1 + |2^{-j} y|)^\alpha dy \\ &\leq CM(j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty}, \end{aligned}$$

where we applied (2.5) and a change of variables to get the second-to-last inequality and (3.7) to get the last inequality above.

For $g_z Q_j^4(u_1, w)$, we again follow [18] and write $Q_j^4(u_1, w)$ as the sum of two terms:

$$Q_j^4(u_1, w) = Q_j^{4,1}(u_1, w) + Q_j^{4,2}(u_1, w),$$

where

$$\begin{aligned} Q_j^{4,1}(u_1, w) &= \sum_{k=1}^d \Delta_j \partial_k R(u_1^k - S_{j-2} u_1^k, w), \text{ and} \\ Q_j^{4,2}(u_1, w) &= \sum_{k=1}^d \{\Delta_j R(S_{j-2} u_1^k, \partial_k w) - R(S_{j-2} u_1^k, \Delta_j \partial_k w)\}. \end{aligned}$$

By properties of Littlewood Paley operators and Lemma 6,

$$\begin{aligned}
|g_z Q_j^{4,1}(u_1, w)| &= \left| g_z \sum_{k=1}^d \Delta_j \partial_k \sum_{\substack{|\iota-\iota'| \leq 1, \\ \max\{\iota, \iota'\} \geq j-3}} \Delta_\iota (u_1^k - S_{j-2} u_1^k) \Delta_{\iota'} w \right| \\
&\leq C \sum_{k=1}^d \sum_{\substack{|\iota-\iota'| \leq 1, \\ \max\{\iota, \iota'\} \geq j-3}} 2^j \|g_z \Delta_\iota (u_1^k - S_{j-2} u_1^k) \Delta_{\iota'} w\|_{L^\infty} \\
&\leq C \sup_q \|g_z \Delta_q w\|_{L^\infty} \sum_{l \geq j-M_0} 2^{j-l} \|\Delta_l \nabla u_1\|_{L^\infty} \leq CM \sup_q \|g_z \Delta_q w\|_{L^\infty},
\end{aligned}$$

where Lemma 9 was again used to get the last inequality.

The strategy for estimating $Q_j^{4,2}(u_1, w)$ is similar to that used to bound $g_z Q_j^2(u_1, w)$. By properties of Littlewood Paley operators and the divergence-free property of u_1 ,

$$\begin{aligned}
Q_j^{4,2}(u_1, w) &= g_z \sum_{k=1}^d \sum_{l=j-M_0}^{j-1} \sum_{l'=l-1}^{l+1} [\Delta_j, \Delta_l S_{j-2} u_1^k] \Delta_{l'} \partial_k w \\
&= \sum_{k=1}^d \sum_{l=j-M_0}^{j-1} \sum_{l'=l-1}^{l+1} 2^{jd} \int_{\mathbb{R}^d} \phi(2^j y) [\Delta_l S_{j-2} u_1^k(x) - \Delta_l S_{j-2} u_1^k(x-y)] \Delta_{l'} \partial_k w(x-y) dy \\
&= \sum_{k=1}^d \sum_{l=j-M_0}^{j-1} \sum_{l'=l-1}^{l+1} 2^{j(d+1)} \int_{\mathbb{R}^d} (\partial_k \phi)(2^j y) [\Delta_l S_{j-2} u_1^k(x) - \Delta_l S_{j-2} u_1^k(x-y)] \Delta_{l'} w(x-y) dy.
\end{aligned}$$

Hence,

$$\begin{aligned}
|g_z Q_j^{4,2}(u_1, w)| &\leq \sum_{l=j-M_0}^{j-1} \sum_{l'=l-1}^{l+1} 2^{j(d+1)} \int_{\mathbb{R}^d} |\nabla \phi(2^j y)| \|\Delta_l \nabla S_{j-2} u_1\|_{L^\infty} |y| g_z(x) h_z(x-y) \|g_z \Delta_{l'} w\|_{L^\infty} dy \\
&\leq \sum_{l=j-M_0}^{j-1} \sum_{l'=l-1}^{l+1} 2^{j(d+1)} \|\Delta_l \nabla u_1\|_{L^\infty} \|g_z \Delta_{l'} w\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla \phi(2^j y)| |y| (1+|y|)^\alpha dy \\
&\leq C \sum_{l=j-M_0}^{j-1} \sum_{l'=l-1}^{l+1} \|\omega_1\|_{L^\infty} \|g_z \Delta_{l'} w\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla \phi(y)| |y| (1+|2^{-j} y|)^\alpha dy \\
&\leq CM \sup_q \|g_z \Delta_q w\|_{L^\infty},
\end{aligned}$$

where we again used (2.5) and Lemma 9.

We conclude that

$$\|II\|_{L^\infty} \leq CM(j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty}.$$

We now estimate IV . We follow arguments from [17]. Set $\tilde{\phi}_j = \phi_{j-1} + \phi_j + \phi_{j+1}$, so that $\tilde{\Delta}_j = \Delta_{j-1} + \Delta_j + \Delta_{j+1}$. As in [17], write

$$(3.12) \quad \begin{aligned} IV &= -g_z \nabla \Delta_j p = -g_z \nabla (-\Delta)^{-1} \sum_{k,l=1}^d \partial_k \tilde{\Delta}_j \partial_l \Delta_j (w^l u_1^k) \\ &\quad - g_z \nabla (-\Delta)^{-1} \sum_{k,l=-1}^d \partial_l \tilde{\Delta}_j \partial_k \Delta_j (w^k u_2^l) =: V + VI. \end{aligned}$$

For V , observe that

$$V = g_z(x) \sum_{k,l=1}^d \int_{\mathbb{R}^d} \nabla (-\Delta)^{-1} \partial_k \tilde{\phi}_j(y) \partial_l \Delta_j (w^l u_1^k)(x-y) dy,$$

so that

$$(3.13) \quad \begin{aligned} |V| &\leq \sum_{k,l=1}^d \|g_z \partial_l \Delta_j (w^l u_1^k)\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla (-\Delta)^{-1} \partial_k \tilde{\phi}_j(y)| g_z(x) h_z(x-y) dy \\ &\leq \sum_{k,l=1}^d \|g_z \partial_l \Delta_j (w^l u_1^k)\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla (-\Delta)^{-1} \partial_k \tilde{\phi}_j(y)| (1+|y|)^\alpha dy, \end{aligned}$$

where we used (2.5) to get the last inequality. The operator $\nabla (-\Delta)^{-1} \partial_k$ can be expressed as a product of Riesz operators, which commute with dilations. Thus, a change of variables gives

$$(3.14) \quad \begin{aligned} |V| &\leq \sum_{k,l=1}^d \|g_z \partial_l \Delta_j (w^l u_1^k)\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla (-\Delta)^{-1} \partial_k \tilde{\phi}(y)| (1+|2^{-j}y|)^\alpha dy \\ &\leq \sum_{k,l=1}^d \|g_z \partial_l \Delta_j (w^l u_1^k)\|_{L^\infty} \int_{\mathbb{R}^d} |\nabla (-\Delta)^{-1} \partial_k \tilde{\phi}(y)| (1+|y|)^\alpha dy. \end{aligned}$$

Observe that, since $\tilde{\phi}$ has Fourier transform supported away from the origin, Lemma 8 can be applied. Thus the integral in (3.14) is finite, and

$$(3.15) \quad |V| \leq C \sum_{k,l=1}^d \|g_z \partial_l \Delta_j (w^l u_1^k)\|_{L^\infty}.$$

We estimate VI in exactly the same way as V , and we conclude that

$$(3.16) \quad |VI| \leq C \sum_{k,l=1}^d \|g_z \partial_k \Delta_j (w^k u_2^l)\|_{L^\infty}.$$

We can now complete the estimate for V and VI by applying an argument identical to that used to estimate III . This gives

$$\|IV\|_{L^\infty} \leq \|V\|_{L^\infty} + \|VI\|_{L^\infty} \leq CM(j-2) \sup_q \|g_z \Delta_q w\|_{L^\infty}.$$

Viewing (3.3) as a transport equation ($g_z \Delta_j w$ transported by $S_{j-2} u_1$), integrating in time, and applying the estimates for *I*, *II*, *III*, and *IV* above, we conclude that for $3 \leq j \leq p_0$,

$$\|g_z \Delta_j w(t)\|_{L^\infty} \leq \|g_z \Delta_j w^0\|_{L^\infty} + CM \int_0^t \left((j-2) \sup_q \|g_z \Delta_q w(s)\|_{L^\infty} \right) ds,$$

so that

$$(3.17) \quad \sup_{3 \leq j \leq p_0} \|g_z \Delta_j w(t)\|_{L^\infty} \leq \sup_{3 \leq j \leq p_0} \|g_z \Delta_j w^0\|_{L^\infty} + CM \int_0^t \left(p_0 \sup_q \|g_z \Delta_q w(s)\|_{L^\infty} \right) ds.$$

For the case $j > p_0$, note that by Bernstein's lemma and Lemma 9,

$$(3.18) \quad \sup_{j \geq p_0} \|g_z \Delta_j w(t)\|_{L^\infty} \leq \sup_{j \geq p_0} 2^{-j} \|\Delta_j \nabla w(t)\|_{L^\infty} \leq CM 2^{-p_0}.$$

Now assume $-1 \leq j \leq 2$. In this case, $\|I\|_{L^\infty}$ can be handled exactly as in (3.4). For *III*, the divergence-free assumption on u_2 and Lemma 6 imply that

$$|III| = |g_z \Delta_j \nabla \cdot (w u_2)| \leq C 2^j \|g_z w u_2\|_{L^\infty} \leq CM \|g_z w\|_{L^\infty}.$$

For *IV*, we write

$$(3.19) \quad g_z \nabla \Delta_j p = g_z \sum_{i,k=1}^d R_i R_k \nabla \Delta_j (w^i u_1^k + u_2^i w^k).$$

We focus on $g_z R_i R_k \nabla \Delta_j (w^i u_1^k)$. Note that for $0 \leq j \leq 2$,

$$g_z R_i R_k \nabla \Delta_j (w^i u_1^k) = \int_{\mathbb{R}^d} (R_i R_k \nabla \phi_j(y)) g_z(x) h_z(x-y) u_1^k(x-y) (g_z w^i)(x-y) dy,$$

so

$$(3.20) \quad \begin{aligned} |g_z R_i R_k \nabla \Delta_j (w^i u_1^k)| &\leq C \|u_1\|_{L^\infty} \|g_z w\|_{L^\infty} \int_{\mathbb{R}^d} |R_i R_k \nabla \phi_j(y)| (1+|y|)^\alpha dy \\ &\leq CM \|g_z w\|_{L^\infty} \int_{\mathbb{R}^d} |R_i R_k \nabla \phi(y)| (1+|2^{-j}y|)^\alpha dy, \end{aligned}$$

where we used (2.5) to get the first inequality. Since $\nabla \phi$ has Fourier support away from the origin, Lemma 8 can be applied to conclude that the integral in (3.20) is finite. Thus,

$$|g_z R_i R_k \nabla \Delta_j (w^i u_1^k)| \leq CM \|g_z w\|_{L^\infty}.$$

For $j = -1$, we must estimate

$$(3.21) \quad g_z R_i R_k \nabla \Delta_{-1} (w^i u_1^k) = g_z(x) \int_{\mathbb{R}^d} R_i R_k \nabla \chi(y) (w^i u_1^k)(x-y) dy.$$

This term is the only term which places a substantial restriction on the growth of h_z (or decay of g_z) at infinity. Indeed, estimates for all other terms in the proof of Theorem 1 will hold with $h_z(x) = (1+|x-z|)^\alpha$ for any $\alpha > 0$. It is in the estimate for (3.21) where we utilize the membership of u_1 and u_2 in $L^p(\mathbb{R}^d)$ to optimize the value of α .

Let q satisfy $1/p + 1/q = 1$. Then

$$g_z R_i R_k \nabla \Delta_{-1} (w^i u_1^k) = g_z(x) \int_{\mathbb{R}^d} R_i R_k \nabla \chi(y) h_z(x-y) u_1^k(x-y) (g_z w^i)(x-y) dy,$$

and hence,

$$(3.22) \quad \begin{aligned} |g_z R_i R_k \nabla \Delta_{-1}(w^i u_1^k)| &\leq \|R_i R_k \nabla \chi(\cdot) g_z(x) h_z(x - \cdot)\|_{L^q} \|u_1^k(g_z w^i)\|_{L^p} \\ &\leq \|R_i R_k \nabla \chi(\cdot)(1 + |\cdot|)^\alpha\|_{L^q} \|u_1\|_{L^p} \|g_z w\|_{L^\infty}, \end{aligned}$$

where we used (2.5) to get the last inequality.

Given p , we must determine values of α for which $\|R_i R_k \nabla \chi(\cdot)(1 + |\cdot|)^\alpha\|_{L^q}$ is finite. In view of Lemma 8, a short calculation shows that $\|R_i R_k \nabla \chi(\cdot)(1 + |\cdot|)^\alpha\|_{L^q}$ is finite if

$$\alpha < 1 + d/p.$$

Therefore, assuming $\alpha < 1 + d/p$, it follows that

$$|g_z R_i R_k \nabla \Delta_{-1}(w^i u_1^k)| \leq CM \|g_z w\|_{L^\infty}.$$

Of course, arguments identical to those above imply that

$$\begin{aligned} |g_z R_i R_k \nabla \Delta_{-1}(u_2^i w^k)| &\leq CM \|g_z w\|_{L^\infty}, \text{ and} \\ |g_z R_i R_k \nabla \Delta_j(u_2^i w^k)| &\leq CM \|g_z w\|_{L^\infty} \end{aligned}$$

for $0 \leq j \leq 2$. Inserting these estimates into (3.19) gives the following estimate for $-1 \leq j \leq 2$:

$$|g_z \nabla \Delta_j p| \leq CM \|g_z w\|_{L^\infty}.$$

We now turn our attention to *II*. Instead of considering the commutator, we estimate each term in the difference individually. For the first term in the difference, we observe that, by Lemma 6,

$$\begin{aligned} |g_z S_{j-2} u_1 \cdot \nabla \Delta_j w| &\leq \|S_{j-2} u_1 g_z \nabla \Delta_j w\|_{L^\infty} \\ &\leq \|S_{j-2} u_1\|_{L^\infty} \|g_z \nabla \Delta_j w\|_{L^\infty} \leq CM 2^j \|g_z \Delta_j w\|_{L^\infty} \end{aligned}$$

for $-1 \leq j \leq 2$. For the second term in the difference, note that by the divergence-free property of u_1 and Lemma 6,

$$|g_z \Delta_j(u_1 \cdot \nabla w)| \leq C 2^j \|u_1 g_z w\|_{L^\infty} \leq CM \|g_z w\|_{L^\infty}.$$

We also recall the following estimate, which is derived in (3.1) and which holds for any $p_0 \geq 0$:

$$(3.23) \quad \|g_z w\|_{L^\infty} \leq C p_0 \sup_q \|g_z \Delta_q w\|_{L^\infty} + CM 2^{-p_0}.$$

Integrating (3.3) in time, applying the above estimates for *I*, *II*, *III*, and *IV* when $-1 \leq j \leq 2$, applying (3.23), and taking the supremum over $-1 \leq j \leq 2$ gives

$$\sup_{-1 \leq j \leq 2} \|g_z \Delta_j w(t)\|_{L^\infty} \leq \sup_{-1 \leq j \leq 2} \|g_z \Delta_j w^0\|_{L^\infty} + CM \int_0^t (p_0 \sup_q \|g_z \Delta_q w(s)\|_{L^\infty} + 2^{-p_0}) ds.$$

Combining this estimate with (3.17) and (3.18) yields

$$\begin{aligned}
(3.24) \quad & \sup_q \|g_z \Delta_q w(t)\|_{L^\infty} \\
& \leq \sup_q \|g_z \Delta_q w^0\|_{L^\infty} + CM \int_0^t \left(p_0 \sup_q \|g_z \Delta_q w(s)\|_{L^\infty} + 2^{-p_0} \right) ds \\
& \leq C\epsilon_0 + CM \int_0^t \left(p_0 \sup_q \|g_z \Delta_q w(s)\|_{L^\infty} + 2^{-p_0} \right) ds,
\end{aligned}$$

where

$$\epsilon_0 := \|g_z w^0\|_{L^\infty},$$

and where in the last inequality we applied Lemma 6.

At this point, we could apply Gronwall's estimate to obtain

$$\sup_q \|g_z \Delta_q w(t)\|_{L^\infty} \leq (C\epsilon_0 + CM2^{-p_0}t) e^{CMp_0t}$$

and choose p_0 in a manner so as to minimize the right-hand side. This will not, however, yield a sufficiently tight estimate. Instead, we will choose p_0 in (3.24) to vary with t (this is valid, because the constant, C , in (3.24) does not depend on p_0) in a way that allows us to apply Osgood's Lemma.

Toward this end, first observe that it follows from Lemma 6 that for all $t \geq 0$,

$$\delta(t) := \frac{1}{C_0 M t} \int_0^t \sup_k \|g_z \Delta_k w(s)\|_{L^\infty} ds \leq 1,$$

where C_0 is the constant in (2.3). Now choose

$$p_0 = 3 - \log_2 \delta(t).$$

Note that $p_0 \geq 3$ as required.

Substituting our chosen value for p_0 into (3.24) yields

$$\begin{aligned}
(3.25) \quad & \sup_q \|g_z \Delta_q w(t)\|_{L^\infty} \leq C_0 \epsilon_0 + CMt\delta(t) + CM^2t(3 - \log_2 \delta(t))\delta(t) \\
& \leq C_0 \epsilon_0 + CM^2t\delta(t)(3 - \log_2 \delta(t)) \leq C_0 \epsilon_0 + CM^2t\delta(t)(3 - \log \delta(t)).
\end{aligned}$$

Setting $t = s$ in (3.25) and integrating both sides from 0 to t , then dividing both sides by $C_0 M t$ gives

$$\delta(t) \leq \frac{C_0 \epsilon_0}{C_0 M} + CM \frac{1}{t} \int_0^t s \delta(s) (3 - \log \delta(s)) ds \leq \frac{\epsilon_0}{M} + CM \int_0^t \delta(s) (3 - \log \delta(s)) ds.$$

Since $x \mapsto x(3 - \log x)$ is increasing on $(0, 1]$ and we know that $\delta(s) \in (0, 1]$, we can apply Osgood's Lemma to obtain

$$-\log(3 - \log \delta(t)) + \log \left(3 - \log \left(\frac{\epsilon_0}{M} \right) \right) \leq CMt.$$

Taking the exponential of both sides twice, we conclude that

$$\delta(t) \leq \bar{\delta}(t) := e^3 \left(\frac{\epsilon_0}{M} \right)^{e^{-CMt}}.$$

Now increase the bound in (3.25) to

$$(3.26) \quad \sup_q \|g_z \Delta_q w(t)\|_{L^\infty} \leq C_0 \epsilon_0 + CM^2 t \delta(t) (4 - \log \delta(t))$$

and note that $\bar{\delta}(t) \leq e^3$. Then since $x \mapsto x(4 - \log x)$ is increasing for $x \in (0, e^3]$, we can replace $\delta(t)$ in (3.26) with $\bar{\delta}(t)$. This gives,

$$(3.27) \quad \begin{aligned} \sup_q \|g_z \Delta_q w(t)\|_{L^\infty} &\leq C_0 \epsilon_0 + CM^2 t e^3 \left(\frac{\epsilon_0}{M}\right)^{e^{-CMt}} \left(4 - \log \left(e^3 \left(\frac{\epsilon_0}{M}\right)^{e^{-CMt}}\right)\right) \\ &= C_0 \epsilon_0 + CM^2 t e^3 \left(\frac{\epsilon_0}{M}\right)^{e^{-CMt}} \left(1 - e^{-CMt} \log \left(\frac{\epsilon_0}{M}\right)\right) := a. \end{aligned}$$

By (3.1),

$$\|g_z w\|_{L^\infty} \leq C(Na + M2^{-N}).$$

Define $F(N) := Na + M2^{-N}$. This has a minimum when

$$(3.28) \quad N = N_0 := \log_2(M \log 2/a),$$

where we treat N_0 as though it were an integer. At this (near) minimum, assuming that $0 < a < M \log 2$, which holds for all sufficiently small $\|g_z w^0\|_{L^\infty}$, we have

$$\begin{aligned} F(N_0) &= a \log_2 \frac{M \log 2}{a} + \frac{Ma}{M \log 2} = M \frac{a}{M \log 2} - M \log 2 \frac{a}{M \log 2} \log_2 \frac{a}{M \log 2} \\ &= M(b - (\log 2)b \log_2 b) = Mb(1 - \log b), \end{aligned}$$

where $b := a/(M \log 2) < 1$. (Note that if $a > M \log 2$, we can do no better than setting $N = 0$, which gives the useless bound, $\|g_z w\|_{L^\infty} \leq CM$.)

Hence, when $a < M \log 2$, we have

$$\|g_z w\|_{L^\infty} \leq CMb(1 - \log b),$$

which is the bound stated in Theorem 1.

Acknowledgements. The problem this paper addresses was suggested to the authors by Helena Nussenzveig Lopes and Milton Lopes Filho in the context of Serfati solutions to the 2D Euler equations. The National Science Foundation provided support for EC through grant DMS-1160704 and support for JPK through grant DMS-1212141.

REFERENCES

- [1] D. Ambrose, J. Kelliher, M. Lopes Filho, and H. Nussenzveig Lopes. Serfati solutions to the 2D Euler equations on exterior domains. To appear *Journal of Differential Equations*, 2015. [1](#), [3](#), [4](#), [5](#)
- [2] J.-M. Bony. Calcul symbolique et propagation des singularités pour les équations aux dérivés partielles non linéaires, em *Ann. Sci. Ecole Norm. Sup.* (4), 14(2): 209-246, 1981. [6](#)
- [3] J.-Y. Chemin. *Perfect Incompressible Fluids*, Vol 14 of *Oxford Lecture Series in Mathematics and its Applications*. Clarendon Press, Oxford, 1988. [6](#)
- [4] J.-Y. Chemin. Une facette mathématique de la mécanique des fluides. Preprint Ecole Polytechnique, 1993. [1](#)
- [5] J.-Y. Chemin and N. Lerner. Flot de champs de vecteurs non lipschitziens et équations de Navier-Stokes. *J. Differential Equations*, 121(2): 314-328, 1995. [6](#)
- [6] E. Cozzi. A finite time result for vanishing viscosity in the plane with nondecaying vorticity. *Commun. Math. Sci.*, 8(4): 851-862, 2010. [4](#)

- [7] E. Cozzi. Solutions to the 2D Euler equations with velocity unbounded at infinity. *J. Math. Anal. App.*, 423(1): 144-161, 2015. [1](#), [8](#)
- [8] Thierry Gallay. Infinite energy solutions of the two-dimensional Navier-Stokes equations. To appear Ann. Fac. Sci. Toulouse (arXiv:1411.5156), 2014. [4](#), [5](#)
- [9] Gunther, N. On the motion of fluid in a moving container, *Izvestia Akad. Nauk USSR, Ser. Fiz. Mat.*, 20: 1323-1348, 1503-1532, 1927; 21: 521-526,735-756, 1927; 22: 9-30, 1928. [1](#)
- [10] Kato, T. Nonstationary flows of viscous and ideal fluids in \mathbb{R}^3 . *J. Funct. Anal.* , 9: 296-305, 1972. [1](#)
- [11] J. Kelliher. A characterization at infinity of bounded vorticity, bounded velocity solutions to the 2D Euler equations. To appear Indiana University Mathematics Journal, 2015. [2](#)
- [12] Lichtenstein, Leon. Über einige Existenzprobleme der Hydrodynamik. Zweite Abhandlung. Nicht-homogene, unzusammendrückbare, reibungslose Flüssigkeiten, *Math. Z.*, 26(1): 196-323, 1927; Über einige Existenzprobleme der Hydrodynamik. III. Permanente Bewegungen einer homogenen, inkompressiblen, zähen Flüssigkeit, *Math. Z.*, 28: 387-415, 1928; Über einige Existenzprobleme der Hydrodynamik. IV. Stetigkeitssätze. Eine Begründung der Helmholtz-Kirchhoffschen Theorie geradliniger Wirbelfäden, *Math. Z.*, 32(1): 608-640, 1930. [1](#)
- [13] A. Majda. Vorticity and the mathematical theory of incompressible fluid flow. *Comm. Pure Appl. Math.*, 39: 187-220, 1986. [1](#)
- [14] Marchioro, Carlo and Pulvirenti, Mario. Mathematical theory of incompressible nonviscous fluids. Springer-Verlag, New York, 1994. [5](#)
- [15] P. Serfati. Solutions C^∞ en temps, n-log lipschitz bornées en espace et équation d'Euler, *C.R. Acad. Sci. Paris*, 320 I: 555-558, 1995. [1](#), [2](#), [3](#), [4](#), [5](#)
- [16] Y. Taniuchi. Uniformly local L^p estimates for 2-D vorticity equation and its application to Euler equations with initial vorticity in *bmo*. *Comm. Math. Phys.*, 248(1): 169-186, 2004. [1](#), [8](#)
- [17] Y. Taniuchi, T. Tashiro, and T. Yoneda. On the two-dimensional Euler equations with spatially almost periodic initial data. *J. Math. Fluid Mech.*, 12(4): 594-612, 2010. [1](#), [3](#), [4](#), [5](#), [8](#), [14](#)
- [18] M. Vishik. Incompressible flows of an ideal fluid with vorticity in borderline spaces of Besov type. *Ann. Sci. Ecole Norm. Sup. (4)*, 32(6): 769-812, 1999. [1](#), [3](#), [5](#), [8](#), [11](#), [12](#)
- [19] J. P. Ward, K. N. Chaudhury, and M. Unser. Decay properties of Riesz transforms and steerable wavelets. *SIAM J. Imaging Sci.*, 6: 984-998, 2013. [7](#)
- [20] V. Yudovich. Nonstationary flows of an ideal incompressible fluid. *Zhurnal Vych Matematika.*, 3: 1032-1066, 1963. [1](#)
- [21] V. Yudovich. Uniqueness theorem for the basic non stationary problem in the dynamics of an ideal incompressible fluid. *Math. Res. Lett.*, 2(1): 27-38, 1995. [1](#), [2](#)

DEPARTMENT OF MATHEMATICS, OREGON STATE UNIVERSITY
E-mail address: cozzie@math.oregonstate.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE
E-mail address: kelliher@math.ucr.edu